

Integrable invariant Sobolev metrics on the Abelian extension of the diffeomorphism group of the circle and two-component generalizations of the Camassa-Holm equation

P. A. Kuzmin

*Department of Physics, Moscow State University,
Leninskie gory, Moscow 119992, Russia
e-mail: kuzminp@list.ru*

Abstract

In this note we classify some integrable invariant Sobolev metrics on the Abelian extension of the diffeomorphism group of the circle. We also derive a new two-component generalization of the Camassa-Holm equation. The system obtained appears to be unique bi-Hamiltonian flow on the coadjoint orbit of $\text{Diff}_+(S^1) \ltimes C^\infty(S^1)$ generalizes the Camassa-Holm flow on $\text{Vect}(S^1)^*$.

It is known that the Camassa-Holm equation can be interpreted as the flow on the coadjoint orbit of the diffeomorphism group of the circle. The Camassa-Holm equation was derived in [6, 7, 9]. Two-component generalizations of the equation were introduced in [1, 2, 3, 4, 5], where some properties of the systems were studied. All these systems appear to be bi-Hamiltonian flows on the regular part of the dual space of the Lie algebra $\mathfrak{g} = \text{Vect}(S^1) \ltimes C^\infty(S^1)$.

In this paper we consider a class of the Hamiltonian flows on the space $\mathfrak{g}_{reg}^*$ with quadratic Hamiltonian functions of special form. We are going to classify all the flows those are bi-Hamiltonian relative to a modified Lie-Poisson structure on $\mathfrak{g}_{reg}^*$. We derive a family of two-component completely integrable generalizations of the Camassa-Holm equation.

There is a standard way to construct a pair of compatible Poisson structures and obtain an integrable flow. Let $\mathfrak{g}$ be a Lie algebra, $\mathfrak{g}^*$ be the dual space of $\mathfrak{g}$ (the regular part of the dual space of $\mathfrak{g}$). One can define the canonical Lie-Poisson structure on $\mathfrak{g}^*$ given by

$$\{f, g\}(m) = m([df, dg]) \quad (1)$$

for all smooth $f, g : \mathfrak{g}^* \rightarrow \mathbb{R}$. Each cocycle $\omega \in Z^2(\mathfrak{g})$ defines another Poisson structure on $\mathfrak{g}^*$, which is compatible with the canonical Lie-Poisson structure. This structure is called the modified Lie-Poisson structure.

Let us recall some facts on the Camassa-Holm equation. The canonical Lie-Poisson structure on the regular part of the dual space of $\text{Vect}(S^1)$ is represented by the skew-symmetric operator

$$j_0(m) = m_x + 2m\partial, \quad m \in \text{Vect}(S^1)_{reg}^*.$$

The regular $\mathfrak{g}_{reg}^*$ part of the dual space is the set of all linear functionals on $\mathfrak{g} = \text{Vect}(S^1)$ of the form

$$\bar{m} : f \mapsto \int_{S^1} f m dx, \quad m \in C^\infty(S^1),$$

and we identify the functional $\bar{m}$ with the function m . [11] For the modified Lie-Poisson structure, which is obtained from the Virasoro cocycle, one has

$$j(m) = 2m_0\partial + (m_0)_x - c_1\partial^3,$$

where $m_0 \in \text{Vect}(S^1)^*$, $c_1 \in \mathbb{R}$, and $\partial = \frac{d}{dx}$. Consider the Hamiltonian vector field $x(m) = j_0(m)\delta h_0(m)$ on $\text{Vect}(S^1)_{reg}^*$ with Hamiltonian function

$$h_0(m) = \frac{1}{2} \int_{S^1} m(1 - \partial^2)^{-1} m dx.$$

Here $\delta\phi(m)$ denotes the gradient of the smooth functional $\phi(m)$. The field $x(m)$ is known to be Hamiltonian with respect to the Poisson structure $j_1(m) = \partial - \partial^3$ with Hamiltonian function

$$h_1(m) = \frac{1}{2} \int_{S^1} (u^3 + uu_x^2) dx, \quad u = (1 - \partial^2)^{-1} m,$$

i.e. $x(m) = j_0(m)\delta h_0(m) = j_1(m)\delta h_1(m)$. The corresponding evolution equation $m_t + x(m) = 0$ is called the Camassa-Holm equation:

$$m_t + m_x u + 2m u_x = 0, \quad u = (1 - \partial^2)^{-1} m.$$

The main goal of this note is to construct analogous bi-Hamiltonian systems on the space $\mathfrak{g}_{reg}^*$, where $\mathfrak{g} = \text{Vect}(S^1) \ltimes C^\infty(S^1)$, and $C^\infty(S^1)$ is the space of all (smooth) tensor densities on S^1 of degree 0, i.e. the $\text{Vect}(S^1)$ -module structure on $C^\infty(S^1)$ is defined as

$$L_f : C^\infty(S^1) \rightarrow C^\infty(S^1), \quad a(x) \mapsto f(x)a'(x),$$

where $f \in \text{Vect}(S^1)$. The canonical Lie-Poisson structure on $\mathfrak{g}_{reg}^*$ is given by the operator

$$J_0 \left(\begin{pmatrix} m \\ p \end{pmatrix} \right) = \begin{pmatrix} m_x + 2m\partial & p\partial \\ p_x + p\partial & 0 \end{pmatrix}, \quad \begin{pmatrix} m \\ p \end{pmatrix} \in \mathfrak{g}_{reg}^*, \quad (2)$$

which defines the coadjoint action of the semi-direct product $\text{Vect}(S^1) \ltimes C^\infty(S^1)$. [12] It is known that $H^2(\text{Vect}(S^1) \ltimes C^\infty(S^1)) = \mathbb{R}^3$ (see, for example, [10, 8]), so the modified Lie-Poisson structure is represented by the constant (i.e. independent on the point $\begin{pmatrix} m \\ p \end{pmatrix} \in \mathfrak{g}_{reg}^*$) operator

$$J = \begin{pmatrix} (m_0)_x + 2m_0\partial - c_1\partial^3 & p_0\partial + c_2\partial^2 \\ (p_0)_x + p_0\partial - c_2\partial^2 & 2c_3\partial \end{pmatrix}, \quad (3)$$

where $\begin{pmatrix} m_0 \\ p_0 \end{pmatrix} \in \mathfrak{g}_{reg}^*$ and c_1, c_2, c_3 are constants. Note that the structures (2), (1) are actually defined on the coadjoint orbit $(\text{Diff}_+(S^1) \ltimes C^\infty(S^1)) / (S^1 \times S^1)$ containing a constant moment $\begin{pmatrix} m_0 \\ p_0 \end{pmatrix} \in \mathfrak{g}_{reg}^*$, $m_0(x) = \text{const}$, $p_0(x) = \text{const}$ [11].

Consider a class of quadratic Hamiltonian functions on $\mathfrak{g}_{reg}^*$ given by

$$H_0 \left(\begin{pmatrix} m \\ p \end{pmatrix} \right) = \frac{1}{2} \left\langle \begin{pmatrix} m \\ p \end{pmatrix}, M^{-1} \begin{pmatrix} m \\ p \end{pmatrix} \right\rangle, \quad (4)$$

where $\langle, \rangle$ denotes the L^2 pairing

$$\left\langle \begin{pmatrix} m_1 \\ p_1 \end{pmatrix}, \begin{pmatrix} m_2 \\ p_2 \end{pmatrix} \right\rangle = \int_{S^1} (m_1(x)m_2(x) + p_1(x)p_2(x)) dx,$$

and the symmetric "inertia" operator is chosen to be of the form

$$M = \begin{pmatrix} \varepsilon(1 - \partial^2) + \sum_{i=2}^{n_0} a_i \partial^{2i} & \sum_{i=0}^{n_1} (-1)^i b_i \partial^i \\ \sum_{i=0}^{n_1} b_i \partial^i & \gamma \end{pmatrix}, \quad (5)$$

where $a_i, b_i, \gamma, \varepsilon$ are constants such that M is strictly positive and invertible. This Hamiltonian function can be considered as a natural generalization of the Hamiltonian function $h_0(m)$. Now we state the main theorem of this note:

Theorem 1 *Assume that m_0, p_0 (see (3)) are constant functions. The only vector field of the form $X = J_0 \delta H_0$ (where J_0, H_0 are given by (2), (4)) that is bi-Hamiltonian relative to the modified Lie-Poisson structure (3) is the Hamiltonian vector field defined by the following "inertia" operator:*

$$M = \begin{pmatrix} \varepsilon(1 - \partial^2) & \alpha - \beta \partial \\ \alpha + \beta \partial & \gamma \end{pmatrix}, \quad (6)$$

where $\alpha, \beta, \gamma, \varepsilon$ are constants.

We obtain the second Poisson structure

$$J_1 = \begin{pmatrix} \varepsilon(\partial - \partial^3) & \alpha \partial - \beta \partial^2 \\ \alpha \partial + \beta \partial^2 & \gamma \partial \end{pmatrix}, \quad (7)$$

and the second Hamiltonian function

$$H_1 \left(\begin{pmatrix} m \\ p \end{pmatrix} \right) = \frac{1}{2} \int_{S^1} (2\alpha u^2 v + 2\beta u u_x v + \gamma u v^2 + \varepsilon u^3 + \varepsilon u u_x^2) dx,$$

$$\begin{pmatrix} u \\ v \end{pmatrix} = M^{-1} \begin{pmatrix} m \\ p \end{pmatrix},$$

such that $X = J_0 \delta H_0 = J_1 \delta H_1$. The equations obtained have the form

$$\begin{cases} m_t + m_x u + 2m u_x + p v_x = 0 \\ p_t + (p u)_x = 0, \end{cases}$$

where $m = \varepsilon u - \varepsilon u_{xx} + \alpha v - \beta v_x$, $p = \alpha u + \beta u_x + \gamma v$.

The proof of the theorem is based on the following simple observation. This method was developed in [13, 14]. Let X be a smooth vector field on $\mathfrak{g}_{reg}^*$. Suppose that X is Hamiltonian with respect to the constant (i.e. independent on the point of $\mathfrak{g}_{reg}^*$) Poisson structure J . This means that there is exists a smooth function $H : \mathfrak{g}_{reg}^* \rightarrow \mathbb{R}$ such that

$$X = J\delta H(m).$$

The operator $d\delta H(m)$ is symmetric, since $H(m)$ is a smooth function. Therefore, the operator

$$S(m) = J(d\delta H(m))J = d(J\delta H(m))J = (dX)J$$

is symmetric. We obtained a necessary condition for X to be Hamiltonian with respect to the Poisson structure J .

We restrict our considerations to the modified Lie-Poisson structures (3) such that m_0, p_0 are constants. Thus we obtain the following condition for X to be Hamiltonian with respect to J :

$$\text{The operator } (dX)J = \begin{pmatrix} u\partial & v_x \\ 0 & u\partial + u_x \end{pmatrix} J + \begin{pmatrix} m_x + 2m\partial & p\partial \\ p\partial + p_x & 0 \end{pmatrix} M^{-1}J \text{ is symmetric,} \quad (8)$$

where

$$J = \begin{pmatrix} 2m_0\partial - c_1\partial^3 & p_0\partial + c_2\partial^2 \\ p_0\partial - c_2\partial^2 & 2c_3\partial \end{pmatrix}, \quad \begin{pmatrix} u \\ v \end{pmatrix} = M^{-1} \begin{pmatrix} m \\ p \end{pmatrix}.$$

Note that the (operator) expression

$$((dX)J) \left(\begin{pmatrix} m \\ p \end{pmatrix} \right) = \begin{pmatrix} u\partial & v_x \\ 0 & u\partial + u_x \end{pmatrix} J + \begin{pmatrix} m_x + 2m\partial & p\partial \\ p\partial + p_x & 0 \end{pmatrix} M^{-1}J \quad (9)$$

is linear in $\begin{pmatrix} m \\ p \end{pmatrix}$. We are now in a position to obtain a polynomial representation for the following trilinear functionals:

$$\left\langle \begin{pmatrix} m_1 \\ p_1 \end{pmatrix}, ((dX)J) \left(\begin{pmatrix} m \\ p \end{pmatrix} \right) \begin{pmatrix} m_2 \\ p_2 \end{pmatrix} \right\rangle \text{ and } \left\langle ((dX)J) \left(\begin{pmatrix} m \\ p \end{pmatrix} \right) \begin{pmatrix} m_1 \\ p_1 \end{pmatrix}, \begin{pmatrix} m_2 \\ p_2 \end{pmatrix} \right\rangle. \quad (10)$$

We can achieve this substituting

$$\begin{pmatrix} m \\ p \end{pmatrix} = \begin{pmatrix} e^{ikx} \\ e^{ilx} \end{pmatrix}, \quad \begin{pmatrix} m_1 \\ p_1 \end{pmatrix} = \begin{pmatrix} e^{ik_1x} \\ e^{il_1x} \end{pmatrix}, \quad \begin{pmatrix} m_2 \\ p_2 \end{pmatrix} = \begin{pmatrix} e^{ik_2x} \\ e^{il_2x} \end{pmatrix}.$$

One can see that expressions (10) vanish unless $l = k$, $l_1 = k_1$, $l_2 = k_2$, $k = -k_1 - k_2$. Therefore

$$\begin{aligned} & \left\langle \begin{pmatrix} e^{ik_1x} \\ e^{ik_1x} \end{pmatrix}, ((dX)J) \begin{pmatrix} e^{-ik_1x-ik_2x} \\ e^{-ik_1x-ik_2x} \end{pmatrix} \begin{pmatrix} e^{ik_2x} \\ e^{ik_2x} \end{pmatrix} \right\rangle, \\ & \left\langle ((dX)J) \begin{pmatrix} e^{-ik_1x-ik_2x} \\ e^{-ik_1x-ik_2x} \end{pmatrix} \begin{pmatrix} e^{ik_1x} \\ e^{ik_1x} \end{pmatrix}, \begin{pmatrix} e^{ik_2x} \\ e^{ik_2x} \end{pmatrix} \right\rangle \end{aligned} \quad (11)$$

are polynomials in k_1 and k_2 . Straightforward calculation yields the result posed: the polynomials (11) coincide and (8) is satisfied only if the "inertia" operator (5) has the form (6). The second Poisson structure turns to be of the form (7).

Let us consider some special cases of the bi-Hamiltonian flow constructed. Setting $\alpha = \beta = 0$, $\varepsilon = -\gamma = 1$ we recover the system studied in [5, 3]:

$$\begin{cases} m_t + m_x u + 2mu_x - pp_x = 0 \\ p_t + (pu)_x = 0, \end{cases} \quad (12)$$

where $m = u - u_{xx}$. If, in turn, we set $\varepsilon = \gamma = 0$, $\alpha = \beta = 1$, we recover the system introduced in [1, 4]:

$$\begin{cases} (v - v_x)_t + (2uv - v_x u)_x = 0 \\ (u + u_x)_t + (u^2 + uu_x)_x = 0, \end{cases} \quad (13)$$

which is equivalent to the system considered in [4] under $x \mapsto -x$, $u \mapsto v$, $v \mapsto u$. The same bi-Hamiltonian structures for (12), (13) were obtained in [5, 3, 1, 4].

References

- [1] Liu S-Q., Zhang Y., Deformations of semisimple bihamiltonian structures of hydrodynamic type, J. Geom. Phys., 2005, vol. 54, pp. 427–453.
- [2] Oliver P.J., Rozenau P, Tri-Hamiltonian duality between solitons and solitary waves having compact support, Phys. Rev. E 53, 1996, pp. 1900–1906.
- [3] Aratyn H., Gomes J.F., Zimerman A.H., On negative flows of the AKNS hierarchy and a class of deformations of a bihamiltonian structure of hydrodynamic type, J. Phys. A: Math. Gen., 2006, vol. 39, 1099–1114.

- [4] Falqui G., On a Camassa-Holm type equation with two dependent variables, *J. Phys. A: Math. Gen.*, 2006, vol. 39, 327–342.
- [5] Ming Chen, Si-Qi Liu, Youjin Zhang, A two-component generalization of the Camassa-Holm equation and its solutions, *Lett. Math. Phys.*, 2006, vol. 75, no. 1, pp. 1–15.
- [6] Camassa R., Holm D., An integrable shallow water equation with peaked solitons, *Phys. Rev. Lett.*, 1993, vol. 71, no.11, pp. 1661–1664.
- [7] Camassa R., Holm D., Hyman J.M., A new integrable shallow water equation, *Advances in Applied Mechanics*, 1994, vol. 31, pp. 1–33.
- [8] Arbarello E., De Concini C., Kac V.G., Procesi C. Moduli, spaces of curves and representation theory, *Commun. Math. Phys.*, 1988, vol. 117, pp. 1–36.
- [9] Fuchssteiner B., Fokas, A.S., Symplectic structures, their Backlund transformations and hereditary symmetries, *Physica D* 4, 1981, pp. 47–66.
- [10] Ovsienko V.Yu., Roger C., Extension of Virasoro group and Virasoro algebra by modules of tensor densities on S^1 , *Funct. Anal. Appl.*, 1996, vol. 31, no.4
- [11] Kirillov A.A., Infinite dimensional Lie groups: Their orbits, invariants and representations. The geometry of moments. *Lect. Notes in Math.* **970**, Berlin–Heidelberg–New York: Springer-Verlag, 1982, pp. 101–123.
- [12] Marcel P., Ovsienko V., Roger C., Extensions of the Virasoro and Neveu-Schwarz Algebras and Generalized Sturm-Liouville Operators, *Lett. Math. Phys.*, 1997, vol. 40, pp. 31–39.
- [13] Constantin A., Kolev B., Integrability of Invariant Metrics on the Diffeomorphism Group of the Circle, *J. Nonlinear Sci.*, 2006, vol. 16, pp. 109–122.
- [14] Constantin A., Kolev B., Lenells J., Integrability of invariant metrics on the Virasoro group, *Phys. Lett. A*, 2006, vol. 360, pp. 75–80.